















EMR Quick Reference Guide

Medications - Warfarin – Prescribing Initial Therapy (For commencing therapy in Warfarin-naive patients)

1. Click on **Orders** and click **+ Add**
2. Type Warfarin in the 'Search' field.
3. Select **Warfarin Therapy Initiation** order set (indicated by )
4. Review the pre-checked orders and select any other orders as necessary. Complete the mandatory fields for any orders with  icon



Warfarin Therapy Initiation (Initiated Pending)

-  Education Warfarin **1**
Please supply and outline Warfarin Booklet to patient as part of education process.
- Medications**
 - Please review Warfarin dosing parameters for patients with Synthetic valves in situ
 - BRIDGING ANTICOAGULATION**
 -  Warfarin Bridging. Review when INR 2.0, Cease when INR 2.0 or greater
 -  enoxaparin **2** Select an order sentence
 - COUMADIN**
 -  For NEW warfarin therapy: Coumadin is the brand of choice
 -  warfarin (Coumadin) **3** Select an order sentence
 - MAREVAN**
 -  warfarin (Marevan) **3** Select an order sentence
 - warfarin dose check
 -  The 'included' warfarin dose check task ensures nursing staff check that there is a warfarin order on 4th day **4**
 -  warfarin dose check **4** 1 dose check, ONCE only, First dose T+3;1400
- Pathology**
 -  International Normalised Ratio (INR) **5**
AM Path Rnd, Spec Type: Blood, T+3;0630, Anticoagulant: Warfarin
Commencing Warfarin therapy

	<u>Education Warfarin</u>	A shared nursing/pharmacy task order to provide warfarin education to the patient
2	<u>Bridging Anticoagulation</u>	Bridging enoxaparin can be ordered if required
3	<u>Warfarin</u>	Coumadin® is pre-ticked as it is Western Health's preferred brand. Select the drop down arrow to select an appropriate sentence based on the indication and corresponding target INR range.
4	<u>Warfarin Dose Check</u>	The warfarin dose check order is placed to task nurses to check if a new warfarin dose order has been placed.
5	<u>INR pathology order</u>	Defaults to order INR on the 4 th day (on phlebotomy's AM path round)

5. Check the **Order Details** for the Warfarin Order

Initiation of warfarin is prescribed with a specified duration of 3 days. After this time, it will drop off the MAR. Therefore there is a "Warfarin Dose Check" order for the nurse to ensure warfarin orders have been updated daily. **To continue warfarin after the three days, order ongoing doses using the WARFARIN THERAPY ONGOING orderset.**

Warfarin dose administration time will default to the next 1600 hrs. If the order is placed AFTER 1600, change the 'Priority' field to NOW. This will schedule the day's dose and continue at 1600 daily thereafter.

- Click on **Orders For Signature** and enter 'Current Clinical History'
- Click on **Sign** and enter your password.
- Check the **MAR** to ensure order(s) are correct.

Current Clinical History (Mandatory)
 Atrial Fibrillation

General Clinical History (Optional)

Scheduled	
warfarin (Coumadin) 5 mg, Oral, Tablet, daily for 3 day(s), First dose 19/10/2018 16:00:00, Stop date 22/10/2018 15:59:00, INR Target: 2.0 - 3.0, Indication: Atrial fibrillation For commencement of therapy warfarin INR	5 mg Not given within 5 days.
warfarin dose check 1 dose check, ONCE only, First dose 22/10/2018 14:00:00, Stop date 22/10/2018 14:00:00 warfarin dose check INR	