



EMR Quick Reference Guide

Infusions – Completing a Bag By Bag Infusion – Nursing/Midwifery

This process starts in the **MAR**.

When the total number of bags in the order has been infused:

1. Disconnect infusion from the patient.
2. Click on Pending tile to document final volume infused
3. Locate the correct infusion under Continuous Infusions in the **MAR**
4. Right click on the **End Bag Tile**
5. Select **Complete Order** and click Yes to address the confirmation alert.

sodium chloride 0.9%
sodium chloride 0.9% infusion (BAG BY BAG) 1,000 mL
1,000 mL, IV Infusion, Rate: 125 mL/hr, 3 bag(s), First dose 03/12/2020 17:27:00,
Stop date 04/12/2020 17:26:00, Total volume (mL): 1,000
Administration Information
Sodium Chloride 0.9%

End Bag
3 of 3

Pending
Last bag started:
04/12/2020 08:56 AEDT

Start New Bag
Complete Order

Complete Order

You have selected to complete the order. Do you wish to continue?

Yes No

6. Refresh the MAR and the order is now greyed out and completed under the Discontinued Continuous Infusions

Discontinued Continuous Infusions

heparin (additive) 25,000 unit(s) [1,430 units/hr]
Sodium Chloride 0.9% infusion 250 mL
250 mL, IV Infusion, Rate: 14.3 mL/hr, First dose 13/11/2020 12:40:00, Use premix
heparin bags, Total volume (mL): 250
Administration Information
heparin

Sodium Chloride 0.9%
sodium chloride 0.9% infusion (BAG BY BAG) 1,000 mL
1,000 mL, IV Infusion, Rate: 125 mL/hr, 3 bag(s), First dose 03/12/2020 17:27:00,
Stop date 04/12/2020 17:26:00, Total volume (mL): 1,000
Administration Information
Sodium Chloride 0.9%

Note: If verbal communication is received from Medical Officer to cease the bag by bag infusion before the total number of bags in the order are completed i.e. at end of bag 2 of 3, the Medical Officer must discontinue the infusion order and final volume infused will be via the FBC as per for Truly continuous infusions.