



EMR Quick Reference Guide

Infusions – Completing a Truly Continuous Infusion and documenting Final Volume Infused – Nursing/Midwifery

1. This starts from the **MAR**
2. When the infusion has been discontinued by the Medical Officer, the infusion will appear **Greyed Out** on the MAR and sits under **Discontinued Continuous Infusion**
3. Disconnect infusion from the patient.

Discontinued Continuous Infusions	
heparin (additive) 25,000 unit(s) [1,430 units/hr]	
Sodium Chloride 0.9% infusion 250 mL	
250 mL, IV Infusion, Rate: 14.3 mL/hr, First dose 13/11/2020 12:40:00, Use premix heparin bags, Total volume (mL): 250	
Administration Information	
heparin	
Sodium Chloride 0.9%	

4. Navigate to **Interactive View** from TOC
5. Click on the **FBC** band and **Refresh** the page
6. Locate the greyed out discontinued order.
7. **Right click** on the current date and time cell and Select **insert time and date**.

Intake Total	
heparin (additive) 25,000 unit(s) [1,430 units/hr] + Sodium Chloride 0.9% infusion 250 mL	1500
6 heparin (additive) 25,000 unit(s) [1,500 units/hr] + Sodium Chloride 0.9% infusion 250 mL	250
sodium chloride 0.9% infusion (BAG BY BAG) 1,000 mL	1500

8. A new column will appear.
9. Enter in the final volume infused
10. **Sign** with the **Green** tick and **Refresh** the page
11. The final volume is now visible on the FBC and the MAR