# Clinical Care – Dynamic Group Use (Peripheral IV Cannula Documentation




Digital Health Quick Reference Guide

## This Quick Reference Guide will explain how to:


Document a Dynamic Group in iView (using E.g. Peripheral IV Cannula).

#### Part 1: Document line

1. Go to Menu → Interactive View → Adult Lines - Devices → Open the Peripheral IV Dynamic Group.




2. Choose the appropriate fields in the **Dynamic Group** and click **OK**.



- The Line assessment is displayed.
- In the Activity DTA choose Insert new site.


Then complete the rest of the assessment as Appropriate.






# Part 2: Document against the line each shift, and if changes occur

- 1. Go to Menu → Interactive View → Adult Lines Devices → Open the Peripheral IV Dynamic Group.
- 2. Click on the **Activity** DTA and choose the **Assessment** option.
- 3. Document against the appropriate DTAs, then click on **OK**.




## Part 3: Document line Discontinued (line to be removed)

- 1. Go to Menu → Interactive View → Adult Lines Devices → Open the Peripheral IV Dynamic Group.
- 2. Choose the correct cannula (the patient may have more than one cannula).
- 3. Click on the Activity band and choose **Discontinued**.
- Choose the appropriate removal reason and then fill in the rest of the fields.



Right click on the cannula title (grey) bar and choose the Inactivate option.



6. The Dynamic Group (assessment) will now be greyed out and inactive. It will also be collapsed automatically (but can be expanded to review existing data by clicking on the small black arrow next to the location (grey) bar).

