Medications – Insulin - Supplemental Subcutaneous Insulin (Sliding Scale)



Digital Health Quick Reference Guide

. Go to Orders and Referrals and click +Add

2. Search by the insulin brand (e.g. Novorapid, Humalog, Apidra), or use search term 'insulin supplemental', or 'insulin sliding scale'

3. Select the orderset

4. This is a multi-phase orderset with each brand in its own phase

Note: novorapid is Western Health's preferred supplemental insulin and hence has the default order action: Order now

Humalog and Apidra are set to 'Do not order' by default, this can be changed to 'Order now' as appropriate

Search:	Search: insulin supplemental] 🔍	Advanced Options		▼ Type:
\Lambda Up	🚮 Home	🚖 Favorites	- 🚞 F	olders	🐚 Сору	Folder:
Insulin SUPPLEMENTAL (Sliding Scale - Novorapid, Humalog, Apidra)						

sulin SUPPLEMENTAL (Sliding Scale - Novorapid, Humalog, Apidra)					
Select Visit and Start Time					
Confirm novoRAPID Start Date/Time					
07/09/2019 🚔 💌 0943 🚔 AEST					
onfirm Phase Ac	tion				
onfirm Phase Ac	Start Date/Time	Action			
ionfirm Phase Ac Phase novoRAPID humALOG	tion Start Date/Time 09/07/2019 09:43 AEST	Action Order now Do not order	• •		
confirm Phase Ac Phase novoRAPID humALOG Apidra	tion Start Date/Time 09/07/2019 09:43 AEST	Action Order now Do not order Do not order	• • •		

- 5. To order novorapid 'now', click OK
- 6. Tick the orders you wish to place for the BGL ranges.

N	Novorapid						
	🔆 Novorapid is the Western Health preferred supplemental insulin						
	今 BGL (mmol/L): 0 - 4: Do not order						
	3 BGL (mmol/L): 4.1 - 8						
	🗹 🚯 🗭 insulin aspart (NovoRapid Penfill) 💽 Select an order sentence						
	BGL (mmol/L): 8.1 - 12						
	Image: State of the second	S, PRN for hyperglycaemia					
	BGL (mmol/L): 12.1 - 16						
	 insulin aspart (NovoRapid Penfill) 6 unit(s), Subcutaneous, Cartridge, TDS, PRN BGL (mmol/L): 12.1 - 16 	for hyperglycaemia					
	3 BGL (mmol/L): 16.1 - 20						
	 insulin aspart (NovoRapid Penfill) 8 unit(s), Subcutaneous, Cartridge, TDS, PRN BGL (mmol/L): 16.1 - 20 	for hyperglycaemia					
	BGL (mmol/L): > 20						
	 insulin aspart (NovoRapid Penfill) Insulin aspart (NovoRapid Penfill) Insulin aspart (NovoRapid Penfill) Insulin aspart (NovoRapid Penfill) 	N for hyperglycaemia er					

7. Use the arrows to the select the appropriate dose for the BGL range if required

GL (mmol/L): 8.1 - 12
0 (zero) unit(s), Subcutaneous, Cartridge, TDS, PRN for hyperglycaemia
4 unit(s), Subcutaneous, Cartridge, TDS, PRN for hyperglycaemia
2 unit(s), Subcutaneous, Cartridge, TDS, PRN for hyperglycaemia
6 unit(s), Subcutaneous, Cartridge, TDS, PRN for hyperglycaemia





8. The BGL range included in the order comments can be altered by modifying the order comments

9. Check the MAR to ensure orders are correct. They will be in the 'PRN' section

	Medications		22/10/2018 16:32	
	PRN			
	insulin aspart (NovoRapid Penfill) 0 (zero) unit(s), Subcutaneous, Cartridge, TDS, PRN for hyperglyssemia, First plose 22/10/2018 16:28:00	PRN	0 (zero) unit(s) Not given within 5 days.	
•	BGL (mmol/L): 4.1 - 8			
	insulin aspart			
	Blood Glucose, Capillary - Non Lab			
	Blood Ketones - Non Lab			
	11	PRN	0 (zero) unit(s)	
	insulin aspart (NovoRapid Penfill)		Not given within 5	
	0 (zero) unit(s), Subcutaneous, Cartridge, TDS, PRN for		days.	
	byperglyczemia, First dose 22/10/2018 16:28:00			
	BGL (mmol/L): 8.1 - 12			
	insulin aspart			
The BGL ranges	Blood Glucose, Capillary - Non Lab			
display face-up on the	Blood Ketones - Non Lab		6	
	ेत्र 🛅	PRN	6 unit(s)	
IVIAR	insulin aspart (NovoRapid Penfill)		Not given within 5 days	
	6 unit(s), Subcutaneous, Cartridge, IDS, PRN for hyperglycsemia, Eirst doce 22/10/2018 16:28:00		udys.	
\ ` ▲	BGL (mmol/l): 12.1 - 16			
$\langle \rangle$	unsuun aspart			
	Blood Glucose, Canillary - Non Lab			
	Blood Ketones - Non Lab			
$\langle \rangle$		PRN	8 unit(s)	
	insulin aspart (NovoPanid Penfill)		Not given within 5	
	8 unit(s) Subcutaneous Cartridge TDS PRN for		days.	
$\langle \rangle$	hynerglycaemia Eirst dose 22/10/2018 16:28:00			
	BGL (mmol/L): 16.1 - 20			
	insuin aspart			
	Blood Glucose, Capillary - Non Lab			
	Blood Ketones - Non Lab			
		PRN	10 unit(s)	
	insulin aspart (NovoRapid Penfill)		Not given within 5	
$\langle \rangle$	10 unit(s), Subcutaneous, Cartridge, TDS, PRN for		days.	
	hyperglycomia, First doce 22/10/2018 16:28:00			
	BGL (mmol/L): > 20 and notify Medical Officer			
	insulin aspart			
	Blood Glucose, Capillary - Non Lab			
	Blood Ketones - Non Lab			

