

# Infusions – Commencing Intermittent Infusions – Nursing/Midwifery



## This Quick Reference Guide will explain how to:

Commencing infusions via the Medication Administration Wizard (MAW).

## Infusions – Commencing Intermittent Infusions – Nursing/Midwifery

All infusions must be **commenced** via the **Medication Administration Wizard (MAW)**

1. Check the box next to the infusion you intend to commence
2. Hover over the **mnemonic** column to review last given dose
3. Click the yellow highlighted field under **Result**, the **MAW Administration Window** will open.
4. Review medication order in **Banner**
5. Review the **diluent and diluent volume** – modify if required
6. Enter the **Site of Administration**
7. Enter in **Second Nurse/Midwife details** if required
8. Click **OK**. Second Nurse/Midwife enters password, sign off MAW Administration window
9. Click **refresh** and check the “**additive dose and diluent volume**” is under correct date and time on the MAR. This volume will autopopulate to the Fluid Balance Chart (FBC) under Medications in the **Interactive View and Fluid Balance** tab.

Medications	08/12/2020 22:00	08/12/2020 15:05	08/12/2020 15:04
<b>Scheduled</b>			
flucloxacillin (additive) Sodium Chloride 0.9% infusion 100 mL 1 g, IV Infusion, 6 hourly, Infuse over 30 min(s), Indication: sepsis, First dose 08/12/2020 16:00:00	1 g Last given: 08/12/2020 15:04 AEDT		
flucloxacillin			1 g
Sodium Chloride 0.9%			100 mL