## ICU – Order ICU Resuscitation Procedure Order Set



Digital Health

## This Quick Reference Guide will explain how to:

Place orders retrospectively following a resuscitation or procedure as directed by ICU Medical Officers via the ICU Resuscitation/Procedure Order Set.

## Important – Most bolus medications ordered (vasoactive agents, sedatives and neuromuscular blockers) within this order set will expire 2 hours after the order set is placed.

- This is to give time for the nurses to retrospectively sign off any orders that were given during the resuscitation/procedure.
- Continuous infusions which are ordered will not expire after 2 hours.

## Place ICU Resuscitation/Procedure Order Set

- 1. Click on Medical Officer View from the Table of Contents.
- 2. Navigate to the ICU Quick Orders mPage/tab.
- Navigate to the ICU Order Sets component and click ICU Resuscitation/Procedure.
  - ICU Continuous Renal Replacement Continuous Renal Replacement Therapy (
     ICU Continuous Renal Replacement Continuous Renal Replacement Therapy (
     ICU Impella Admission ICU Impella A
     ICU Resuscitation/Procedure In
     ICU Sepsis ICU Sepsis
- 4. The **shopping cart icon** on the top right will light up and display a 1. Click the icon.
- 5. Click Modify Details.

Modify Details




Medical Officer View

ICU Quick Orders







**Digital Health** 

CTING BEST

Digital Health Quick Reference Guide

🕂 Add   🦨 Document	Medication by Hx   Reconciliation *	Reconciliation Status Meds History  Admission Discharge
Orders Medication Lis	Document In Plan	
	😽 🚫 🕂 Add to Phase - Start: Now Duration: None	
View	♥ Component Status Dose	Details ^
ICU Resusci	ICU Resuscitation/Procedure (Initiated Pending)	
+ Suggested Plans (	ICU Resuscitation/Procedure	0
Orders	() MEDICATIONS	Č.
Non Categoriz	A Resuscitation Medications	0
Communicati	adrenaline (epinephrine) (adrenaline (epinephrine) 1 mg/mL (1:1000) injectable solution)	1 mg, IV Injection, Ampoule, See Instructions, PRN for other: see indication, for 2 hr(s), Indica
- Diet - Patient Care	amiODAROne (amiODAROne for injection)	150 - 450 mg, IV Injection, Ampoule, See Instructions, PRN for other: see indication, for
Continuous In	🗖 😒 atropine	600 MICROg, IV Injection, Ampoule, See Instructions, PRN for other: see indication, for
Pathology	🗖 😒 metaraminol	0.5 mg, IV Injection, Syringe, See Instructions, PRN for other: see indication, for 2 hr(s), Indica
Diagnostic Te:	Iidocaine (lignocaine) (lidocaine (lignocaine) 1% injectable solution)	100 mg, IV Injection, Ampoule, See Instructions, PRN for other: see indication, for 2 hr(s), Indica
Consults	Sedative and Neuromuscular Blocking Agents	0
Referrals	fentanyl (fentanyl for injection)	25 – 200 MICROg, IV Injection, Ampoule, See Instructions, PRN for other: see indication, for
Admit/Transfe ♥		
< >	The Details	
Diagnoses & Problems Related Results	Dx Table Orders For Cosignature Save as My Favorite	Orders For Signature Cancel
/		Sign

- 7. Enter any mandatory fields, then click Sign
- 8. Enter your credentials and click OK.
- 9. Check Orders and Referrals, MAR and MAR Summary to ensure the orders have been placed as intended.